

# Long-Term Effects of Adding Nutrients to an Oligotrophic Coastal Environment

Anna R. Armitage,<sup>1,2\*</sup> Thomas A. Frankovich,<sup>3</sup> and James W. Fourqurean<sup>3</sup>

<sup>1</sup>Department of Biological Sciences and Southeast Environmental Research Center, Florida International University, Miami, Florida 33199, USA; <sup>2</sup>Department of Marine Biology, Texas A&M University at Galveston, PO Box 1675, Galveston, Texas 77553, USA;

<sup>3</sup>Marine Sciences Program, Department of Biological Sciences and Southeast Environmental Research Center, Florida International University, North Miami, Florida 33181, USA

## ABSTRACT

Management of ecological disturbances requires an understanding of the time scale and dynamics of community responses to disturbance events. To characterize long-term seagrass bed responses to nutrient enrichment, we established six study sites in Florida Bay, USA. In 24 plots (0.25 m<sup>2</sup>) at each site, we regularly added nitrogen (N) and phosphorus (P) in a factorial design for 7 years. Five of the six sites exhibited strong P limitation. Over the first 2 years, P enrichment increased *Thalassia testudinum* cover in the three most P-limited sites. After 3 years, *Halodule wrightii* began to colonize many of the P-addition plots, but the degree of colonization was variable among sites, possibly due to differences in the supply of viable propagules. *Thalassia* increased its allocation to aboveground tissue in response to P enrichment; *Halodule* increased in total biomass but did not appear to change its aboveground: belowground tissue

allocation. Nutrient enrichment did not cause macroalgal or epiphytic overgrowth of the seagrass. Nitrogen retention in the study plots was variable but relatively low, whereas phosphorus retention was very high, often exceeding 100% of the P added as fertilizer over the course of our experiments. Phosphorus retentions exceeding 100% may have been facilitated by increases in *Thalassia* aboveground biomass, which promoted the settlement of suspended particulate matter containing phosphorus. Our study demonstrated that low-intensity press disturbance events such as phosphorus enrichment can initiate a slow, ramped successional process that may alter community structure over many years.

**Key words:** aboveground and belowground biomass; epiphyte; eutrophication; macroalgae; nutrient; press; pulse; ramp disturbances; seagrass.

---

Received 11 October 2010; accepted 19 January 2011;  
published online 15 February 2011

**Electronic supplementary material:** The online version of this article (doi:10.1007/s10021-011-9421-2) contains supplementary material, which is available to authorized users.

**Author Contributions:** ARA performed the research, developed methods, analyzed the data, and wrote the paper; TAF performed the research, developed methods, and contributed to writing the paper; JWF conceived of the study, developed methods, and contributed to writing the paper.

\*Corresponding author; e-mail: armitaga@tamug.edu

## INTRODUCTION

Differences among plant species in their ability to compete with other species and utilize resources can dictate community composition and successional patterns after disturbance events (Grime 1977). Disturbance can be broadly defined to include alterations in resources (Pickett and others 1989), such as changes in the availability of limiting nutrients that subsequently alter plant community structure and composition (Fourqurean and others

1995; Siemann and Rogers 2003). These alterations are often caused by anthropogenic nutrient enrichment (Vitousek and others 1997), which can be characterized as pulse (short-term, discrete events), press (sudden events that maintain a constant level), or ramp (events that increase in intensity over time) disturbances depending on the intensity and persistence of the nutrient input (Lake 2000). The response of a system to any of these types of nutrient loading events is strongly dependent on landscape configuration and the availability of propagules (Lotze and others 2000), but in general, nutrient additions increase the biomass and growth rate of plants (Tilman 1987; Micheli 1999). Nutrient enrichment also frequently shifts the competitive balance from slower-growing species that dominate in low nutrient environments to faster-growing species (Duarte 1995; Bargali 1997).

Urbanized and developed shorelines make coastal ecosystems such as seagrass beds vulnerable to anthropogenic impacts. Proximity to urbanized and cultivated terrestrial landscapes exposes seagrass beds to runoff replete with anthropogenic nutrients, which can subsequently alter vital ecological processes and economic functions of seagrass beds (Martinetto and others 2006). Within seagrass beds, nutrient input often increases productivity and plant biomass (Lee and Dunton 2000) and facilitates the overgrowth or replacement of seagrass by fast-growing, opportunistic primary producers (Fourqurean and others 1995; McGlathery 2001).

Nutrient enrichment in seagrass beds may cause changes that extend over many years, especially in oligotrophic systems. For example, in a small-scale pulsed nutrient addition experiment in seagrass beds in Florida Bay, the slow-growing species *Thalassia testudinum* originally responded to fertilization with increases in density and plant size; these increases remained until the fertilized sites were colonized by the faster growing species *Halodule wrightii* (Powell and others 1989; Fourqurean and others 1995). After that colonization event, *Thalassia* biomass and density began to decline, and within 3 years, *Halodule* biomass exceeded that of *Thalassia* (Fourqurean and others 1995). However, primary production in the fertilized plots remained higher than the surrounding area because of the greater mass-specific productivity of *Halodule*, even 20 years after cessation of fertilization (Herbert and Fourqurean 2008).

The fate of nutrients entering the system will have substantial impacts on ecosystem structure and function. If inputs of nitrogen and phosphorus, for example, are retained in the system, then even short-term pulsed or episodic deposition can have

long-term impacts on community characteristics. If those nutrient inputs are quickly lost, then there will be a more rapid return to the base state. Herbert and Fourqurean (2008) found that the consequences of 3 years of fertilization, including increases in biomass, increases in biodiversity, and enhanced primary productivity, persisted for over two decades following the cessation of fertilizer application in Florida Bay. Therefore, it is important to understand the storage of nutrients in the system to predict the long-term impacts of short-term changes in nutrient input.

Previous enrichment experiments performed at these sites have unequivocally demonstrated strong short-term responses (1.5 years) to phosphorus (P) enrichment at sites in eastern Florida Bay, where P addition increased *Thalassia* percent cover and productivity by approximately twofold (Armitage and others 2005). In contrast, *Thalassia* beds in western Florida Bay showed few responses to P addition on that time scale (Armitage and others 2005). However, the expected replacement of *Thalassia* with more opportunistic species like *Halodule* did not occur in that time frame (Powell and others 1989; Fourqurean and others 1995). Furthermore, the storage of the experimental nutrient additions has not yet been quantified. Therefore, the objectives of the present study are to (1) characterize longer-term (7 years) response to experimental nutrient addition within the benthic producer community, and (2) quantify the fate of nutrients added to the system to better predict the long-term responses of seagrass bed structure to the low-intensity press disturbance events of phosphorus enrichment.

## METHODS

### Study Design

To evaluate the long-term effects of N and P enrichment across a P-availability gradient within Everglades National Park in Florida Bay, we used a three-way repeated measures ANOVA design, where the factors were P addition, N addition, and site. The original study concept had two sites nested in each of three major regions of the Bay: Northeast, Interior-Central, and Gulf, as defined by Zieman and others (1989) based on macrophyte and sediment characteristics. However, subsequent analyses revealed that the region designation did not follow expected nutrient-limitation patterns (Armitage and others 2005). Therefore, we considered site to be a factor with six independent levels. In September 2002 we established six study sites (all depths <2 m) in Florida Bay. The three

eastern sites (Duck, South Nest, Bob Allen; see map in online Appendix 1) were characterized by a sparse, short *T. testudinum* canopy with some calcareous green macroalgae, primarily *Penicillus* spp. These three sites occurred in an area of severe P limitation (Fourqurean and Zieman 2002; Armitage and others 2005). Two middle sites (Rabbit Key, Nine Mile Bank) occurred in a region of intermediate P limitation (Fourqurean and Zieman 2002) and featured a dense, tall *T. testudinum* canopy and little macroalgae. The westernmost site (Sprigger Bank) was located in a region that may experience both N and P limitation and was characterized by a dense macroalgal-*Syringodium filiforme* (manatee grass)-*T. testudinum* assemblage. At each site we established twenty-four 0.25-m<sup>2</sup> study plots demarcated with a PVC frame secured to the benthos at one meter intervals.

We randomly assigned treatments (control [C], nitrogen only [N], phosphorus only [P], both nitrogen and phosphorus [NP]) to six plots per site (at Sprigger Bank, replication decreased to three due to the loss of 12 plots from erosion and boat disturbance over the course of the study). N was added in the form of slow release nitrogen fertilizer (Polyon, Pursell Technologies Inc., 38-0-0) and P as granular phosphate rock (Multifos, IMC Global, Ca<sub>3</sub>(PO<sub>4</sub>)<sub>2</sub>). Loading rates of 1.43 g N m<sup>-2</sup> day<sup>-1</sup> and 0.18 g P m<sup>-2</sup> day<sup>-1</sup> (molar N:P ratio 17.6:1) were selected based on potential sewage loading rates (MCSM 2001) and previous studies in the region (Ferdie and Fourqurean 2004). We began bimonthly applications of fertilizer in September 2002 by sprinkling granular fertilizer evenly on the sediment surface and gently working it into the sediment by hand. Sediment in the control plots was similarly disturbed but no fertilizer was added. Benthic fertilizer applications ensured accessibility of nutrients to both aboveground and benthic primary producers (Ferdie and Fourqurean 2004; Mutchler and others 2004). Earlier studies confirmed that this protocol was effective in enriching sediment, seagrass tissue, and seagrass epiphytes (Armitage and others 2005). Bimonthly fertilization continued through April 2006, at which point a portion of each plot was destructively sampled for belowground biomass (see detailed methods below). After a 16-month recovery period, quarterly fertilization was resumed in August 2007 and continued through the end of the study in June 2009.

### Temporal Patterns

Every 3 months from October 2002 to August 2005, we recorded macrophyte percent cover,

epiphyte biomass, benthic microalgal biomass, and seagrass and epiphyte tissue nutrient content. Macrophyte percent cover monitoring continued as frequently as logistics permitted; measurements were taken in September 2007, April 2008, September 2008, December 2008, and June 2009.

We determined macrophyte (seagrass and macroalgae) percent cover using a modified Braun-Blanquet (BB) abundance scale used for extensive monitoring in this region (Fourqurean and others 2001). On this scale, 0 = absent; 0.1 = one individual, less than 5% cover; 0.5 = few individuals, less than 5% cover; 1 = many individuals, less than 5% cover; 2 = 5–25% cover; 3 = 25–50% cover; 4 = 50–75% cover; and 5 = 75–100% cover. Seagrass cover measurements began in October 2002; macroalgal measurements were initiated in February 2004.

We measured seagrass photosynthetic epiphyte biomass by collecting two *T. testudinum* short-shoots from each plot and removing the epiphytes by gently scraping the leaves with a razor blade. Epiphytes were freeze dried and pigments extracted with 90% acetone. The chlorophyll *a* (chl *a*) concentration was determined using a Shimadzu RF-Mini 150 fluorometer with low bandwidth filters (Welschmeyer 1994) following calibration with a chl *a* standard using a Shimadzu UV Mini 1240 spectrophotometer and the spectrophotometric equations of Jeffrey and Humphrey (1975). Epiphyte biomass is reported as µg chl *a* cm<sup>-2</sup> of two-sided seagrass leaf area. Due to logistical constraints, epiphytes were collected quarterly from October 2002 through August 2005 and in April 2006 and April 2008.

We measured the N and P content and calculated the molar N:P ratio of dried seagrass leaves for each of the sampling events through April 2006. N contents were determined using a CHN analyzer (Fisons NA1500). P contents were determined by a dry-oxidation, acid hydrolysis extraction followed by a colorimetric analysis of phosphate concentration of the extract (Fourqurean and others 1992a).

All data collected over time were analyzed with three-way repeated measures ANOVA, where the fixed factors were ±P, ±N, and site. Variances were tested for homoscedasticity with the *F*<sub>max</sub> test and data were log transformed to conform to the assumptions of ANOVA. To compensate for potential violations of sphericity in the data, Greenhouse-Geisser corrections are reported.

### Biomass and Nutrient Budget

To assess macrophyte (seagrass and macroalgae) and epiphyte biomass responses to the enrichment

treatments, we destructively sampled a portion of each plot in April 2006. Using a circular core (15 cm diameter) to subsample biomass in each plot, we removed all aboveground and belowground tissue to a depth of 15 cm. Cores were frozen pending processing in the lab. Plants were separated by species, and for seagrass, tissue was divided into aboveground (photosynthetic) and belowground tissue (rhizomes + roots + other non-photosynthetic tissue). Epiphytes were removed from seagrass leaves by gently scraping with a razorblade. Macroalgal tissue was rinsed to remove adhered sediments but epiphytes were not removed from the complex thalli. Seagrass and macroalgae tissues were dried at 60°C and weighed to determine biomass.

To determine the fate of added nutrients, we recorded the N and P content of each component of the system. Following the collection of biomass cores in April 2006, we measured the N and P content (as described above) of dried seagrass (aboveground and belowground), macroalgal and epiphyte tissues. Sediment cores were not collected in April 2006, so we estimated sediment N and P content using sediment nutrient data from cores (1 cm diameter, 5 cm deep) that were collected from each plot in February 2004. Because these cores were taken earlier in the study, it is likely that we underestimated the sediment nutrient content at the time the plant biomass was recorded.

To determine the long-term importance of nutrient enrichment, we calculated the retention efficiency of nitrogen and phosphorus within 0.25 m<sup>3</sup> compartments (50 × 50 × 100 cm) containing 200 l of water column and 50 l of sediment (sensu Ferdie and Fourqurean 2004). The components included in the calculation were seagrass leaves (separated by species), seagrass belowground tissue (separated by species), macroalgae, *T. testudinum* epiphytes, sediment, and water column. Quantities of nutrients in each component were calculated as nutrient content as a percent of dry weight and biomass (g dry weight). Average values for each treatment at each site were used in the budget calculation. Water column nutrient concentrations from April 2006 were obtained from the Southeast Environmental Research Center Water Quality Monitoring Network at Florida International University. Calculations included net system retention ([total system nutrient (g), enriched] – [total system nutrient (g), control] and retention efficiency ([net system retention (g)/mass applied (g) × 100%).

Following this destructive sampling of biomass, the plots were undisturbed (no sampling or

fertilization) for a period of 16 months, after which the holes created by the core extractions had filled in and were no longer visible, and the seagrass canopies had apparently recovered. The enrichment effects were still visually evident, so we resumed the study in August 2007 and continued with quarterly fertilization and semiannual monitoring of plant percent cover.

Biomass data were analyzed with a three-way fixed factor ANOVA (P addition, N addition, and site) following verification of homoscedasticity with the  $F_{\max}$  test. Biomass data were log transformed if necessary to conform to the assumptions of ANOVA.

## RESULTS

### Temporal Responses

Seagrass leaf nutrient content reflected nutrient treatments, though the magnitude of the responses varied among sites and over time. A highly significant date × site × P interaction (df = 39,  $F = 2.3$ ,  $P < 0.001$ ) was driven by the large decrease in *Thalassia* leaf N:P ratio in P addition treatments at all sites except for the westernmost site, Sprigger Bank (Figure 1, online Appendix 2a). The magnitude of the response to P addition was largest at the three eastern sites and intermediate at Rabbit Key and Nine Mile Bank. A significant 4-way interaction between date, site, P addition, and N addition (df = 31,  $F = 1.5$ ,  $P = 0.031$ ) was driven by higher leaf N:P ratios in NP relative to P only treatments at Duck and Bob Allen Keys on most dates.

The response of *Thalassia* percent cover (as represented by the Braun-Blanquet cover score) varied with site and over time (online Appendix 2b). *Thalassia* cover increased in P addition treatments (P alone and with nitrogen), but only at the three eastern sites (date × site × P df = 52,  $F = 7.6$ ,  $P < 0.001$ ; Figure 2). At those sites, *Thalassia* response to P addition peaked approximately 2 years into the study (September 2004–February 2005), after which *Thalassia* cover declined to control levels in all treatments at most sites. The exception was Bob Allen Key, where *Thalassia* cover decreased but remained measurably higher in P addition plots.

There were no interactions between N and P addition for *Thalassia* cover, but there was a significant date × site × N addition interaction (df = 52,  $F = 1.7$ ,  $P = 0.001$ ; online Appendix 2b). This interaction may have been driven by higher *Thalassia* cover in NP relative to P-only treatments




Fig. 1. Changes in the molar N:P ratio of *Thalassia testudinum* aboveground tissue over time and in response to nutrient (nitrogen, N and phosphorus, P) enrichment. Sites are listed in order from east to west. Bars represent standard error.

at some sites (Duck, Bob Allen) on some dates (May 2004–2005) (Figure 2).

There was a significant interaction between date, site, and P addition for *Halodule* percent cover ( $df = 23$ ,  $F = 4.2$ ,  $P < 0.001$ ; online Appendix 2b). This interaction was driven by a proliferation of *Halodule* in P addition plots starting in the 4th year of the study at all sites except Sprigger Bank (Figure 2). *Halodule* rarely occurred in our control or N only plots. In two of the three eastern sites (Duck and Bob Allen), *Halodule* cover was equivalent to or higher than *Thalassia* cover. There were no significant N effects on *Halodule* cover, and no  $N \times P$  interactions.

A third species of seagrass, *Syringodium filiforme*, was common at the westernmost site, Sprigger

Bank. No statistical analyses were performed on its cover because it occurred only at that site, but qualitative observations suggest that it was consistently more abundant than *Thalassia* (Figure 2f). *Halodule* was not found at Sprigger Bank on any of the sampling dates.

Cover of the two dominant macroalgal groups, calcified green algae and uncalcified red algae, varied over time, though the nature of the temporal changes differed between sites (red macroalgae date  $\times$  site  $df = 7$ ,  $F = 37.6$ ,  $P < 0.001$ ; green macroalgae date  $\times$  site  $df = 7$ ,  $F = 11.7$ ,  $P < 0.001$ ; online Appendix 2c, Figure 3). Overall, macroalgal cover was higher at Sprigger Bank than at all other sites. There were marginally significant site  $\times$  P ( $df = 1$ ,  $F = 4.4$ ,  $P = 0.045$ ) and site  $\times$  N ( $df = 1$ ,






Fig. 2. Changes in *Thalassia testudinum* and *Halodule wrightii* Braun-Blanquet percent cover score over time and in response to nutrient enrichment. There were no significant nitrogen effects and no nitrogen  $\times$  phosphorus interactions; therefore, only control and nitrogen + phosphorus treatments are shown for clarity. Sites are listed in order from east to west. Bars represent standard error.  $\phi$  Note that grey symbols at Sprigger represent *Syringodium filiforme*; *Halodule* was absent at that site.

$F = 4.6$ ,  $P = 0.042$ ) interactions for red macroalgae (online Appendix 2c), but the magnitude of the site and date effects were much larger than any nutrient addition effects. There were no  $N \times P$  interactions for either type of macroalgae.

*Thalassia* epiphyte biomass ( $\mu\text{g chl } a \text{ cm}^{-2}$  seagrass leaf) was lower in P and NP treatments on some dates at Bob Allen (date  $\times$  site  $\times$  P df = 18,  $F = 3.0$ ,  $P < 0.001$ ; online Appendix 2d, Figure 4). Epiphyte biomass was higher at Sprigger Bank than at all other sites, and the magnitude of the site effect was much larger than the temporally variable P-addition effect. Epiphyte biomass did not change in response to nitrogen addition, and there were no  $N \times P$  interactions.

## Biomass

Aboveground *Thalassia* biomass was impacted by P addition, but that effect varied among sites, as revealed by a significant site  $\times$  P interaction (df = 5,  $F = 8.8$ ,  $P < 0.001$ ; online Appendix 2e, Figure 5a). Relative to controls, aboveground *Thalassia* biomass was 2.8–6.7 times higher in P-addition plots at the three eastern bay sites, Duck, South Nest, and Bob Allen. There was no substantial change in aboveground biomass among treatments at the three western sites.

The effect of P addition on belowground *Thalassia* biomass varied significantly among sites (site  $\times$  P df = 5,  $F = 5.2$ ,  $P < 0.001$ ; online Appendix 2e,




Fig. 3. Changes in uncalcified red and calcareous green macroalgae Braun-Blanquet percent cover score over time and in response to nutrient enrichment. There were no significant nitrogen  $\times$  phosphorus interactions; therefore, only control and nitrogen + phosphorus treatments are shown for clarity. Sites are listed in order from east to west. Bars represent standard error.

Figure 5b). Belowground biomass was 1.7 times higher in P addition plots at Bob Allen Keys and about 25% lower in P addition plots at Rabbit and Nine Mile Bank. There was no substantial change in absolute belowground biomass among treatments at the other three sites.

The effect of P addition on the ratio of aboveground to belowground *Thalassia* biomass also varied significantly among sites (site  $\times$  P df = 5,  $F = 4.8$ ,  $P = 0.001$ ; online Appendix 2e, Figure 5c). Relative to controls, the ratio was 3.3–4.5 times higher (that is, more relative aboveground biomass) in P-addition plots at the three eastern bay sites. There was no substantial change in the ratio among treatments at the three western sites.

*Halodule* biomass was patchy and was entirely absent from a large number of plots; precluding statistical analyses. However, *Halodule* biomass was substantially higher in P-addition plots (alone or with nitrogen) at Duck and Bob Allen Keys than in control or nitrogen-only plots (Figure 6). At Rabbit Keys and Nine Mile Bank, *Halodule* biomass was highest in NP addition plots. *Halodule* was absent from Sprigger bank and occurred in only one P-addition plot at South Nest Key.

Macroalgal biomass was highly variable among sites, precluding statistical analyses. Overall, there was no macroalgal proliferation or overgrowth of the seagrasses in enriched plots. Calcareous green algae (*Penicillus* spp. and *Halimeda* spp.) were the




Fig. 4. Changes in *Thalassia testudinum* epiphyte chlorophyll *a* concentration ( $\mu\text{g cm}^{-2}$  of seagrass leaf) over time and in response to nutrient (nitrogen, N and phosphorus, P) enrichment. Sites are listed in order from east to west. Bars represent standard error.

most common genera, with substantially higher biomass at Sprigger Bank than at all other sites (Figure 7). Uncalcified red algae (for example, *Polysiphonia* spp., *Laurencia intricata*, *Palisada poiteau*) were more common in the eastern bay (Figure 7).

### Nutrient Retention Budget

Nitrogen and phosphorus concentrations in seagrass above- and belowground tissue, macroalgal tissue, *Thalassia* epiphytes, soil, and the water column are reported in online Appendix 3. There was little (<31%) nitrogen retention in nitrogen-only plots at most sites, except for one of the western sites, Nine Mile Bank (Table 1). At the three eastern sites, nitrogen retention was higher (24–85%) in plots where both nitrogen and phosphorus were

added. Nearly all of the phosphorus added to the experimental plots was retained in the study plots, particularly within the sediment and belowground *Thalassia* tissue (Table 1). In most cases, phosphorus retention appeared to be greater than 100%, suggesting that the plants or soils were scavenging phosphorus from other sources.

### DISCUSSION

The release from phosphorus limitation caused both acute and longer-term effects on community composition and biomass. Immediate effects of enrichment included an increase in *Thalassia* cover, but these short-term effects did not reflect the long-term, ecosystem level responses. Rather, the most






Fig. 5. *Thalassia testudinum* aboveground and belowground biomass ( $\text{g/m}^2$ ) at six sites after 4 years of nutrient (nitrogen, N and phosphorus, P) enrichment. Bars represent standard error. Note different y axes.

striking outcome of nutrient enrichment in our study was a gradual shift in seagrass species from *Thalassia* to *Halodule*, with a concurrent increase in *Halodule* biomass. As an ecological opportunist with a high potential intrinsic rate of growth, *Halodule* has been shown to overgrow *Thalassia* in nutrient-replete conditions in Florida Bay (Fourqurean and others 1995).

Despite the strong *Halodule* response to P addition, the degree of *Halodule* colonization was




Fig. 6. *Halodule wrightii* aboveground and belowground biomass ( $\text{g/m}^2$ ) at six sites after 4 years of nutrient (nitrogen, N and phosphorus, P) enrichment. Bars represent standard error. Note different y axes.

independent of the severity of phosphorus limitation at each site. Heterogeneity in the supply of viable propagules may have driven the differences in *Halodule* colonization among sites (Lotze and others 2000). One of the most P-limited sites, South Nest Key, had little *Halodule* colonization until the 7th year of the study, whereas the other severely P-limited sites experienced substantial *Halodule* colonization after 3 years of fertilization. In two of the less P-limited sites, Rabbit Key and Nine Mile




Fig. 7. Macroalgal aboveground biomass (g/m<sup>2</sup>) at six sites after 4 years of nutrient (nitrogen, N and phosphorus, P) enrichment. Note different y-axis scale for Sprigger (f).

Bank, *Halodule* colonized a few plots after 1 year of fertilization, but did not spread into other plots during the rest of the study period. This spatial heterogeneity in *Halodule* recruitment, both within and among sites, strongly suggests that the community response to nutrient enrichment is largely influenced by the existing bank of viable seagrass material and the stochastic landings of opportunistic *Halodule* propagules in our plots. Propagule dispersal has been shown to influence species composition in wetland (Campbell and others 2003), marine (Reed and others 2000), and terrestrial (Dosch and others 2007) habitats. The high level of small- (within sites) and large-scale (among sites) spatial heterogeneity in *Halodule* density suggests that the supply of propagules via dispersal is highly variable, even when potential parent plants are established close by (Fourqurean and others 1995).

Overgrowth of fast-growing macro- or epiphytic algae is a commonly reported manifestation of enrichment in seagrass beds (for example, Duarte 1995; Valiela and others 1997; Hauxwell and others 2001; Hughes and others 2004), but we did not detect any algal proliferation in enriched plots despite the high nutrient loading rates and long duration of our experiments. Our results parallel other regional studies of the controls of epiphytic and macroalgal abundance and further suggest that algal loads do not universally increase in seagrass beds during eutrophication. In Florida Bay, nutrient availability explained only a small proportion of the variation in epiphyte loads (Frankovich and Fourqurean 1997), and in a

recent analysis, epiphyte loads were independent of natural gradients in nutrient availability across the seagrass beds of southern Florida (Fourqurean and others 2010). Macroalgal species composition was highly variable among sites (Armitage and others 2005; Armitage and Fourqurean 2009), and this uneven species distribution may have obscured a bay-level macroalgal response to nutrients. Alternatively, nutrient enrichment may have increased the nutritional quality of algal tissue, subsequently stimulating herbivory and effectively preventing algal overgrowth (Heck and others 2006).

*Thalassia* increased its allocation to aboveground biomass in enriched treatments, consistent with fertilization responses observed in other seagrass species (for example, Short 1983; Perez and others 1994; Lee and Dunton 1999). Similar allocation strategies in nutrient-replete conditions have been detected in terrestrial grassland systems (Haase and others 2008; Johnson and others 2008). Our study also agrees with a previous study showing that enrichment does not cause *Halodule* to increase its allocation to aboveground tissue (Dunton 1996), although total *Halodule* biomass in control and nitrogen-only plots was very low, limiting our ability to accurately calculate aboveground:belowground biomass ratios in those treatments.

A substantial portion of the nutrients that were experimentally added to this system were retained in the sediment and belowground seagrass tissue, similar to findings at sites on the ocean side of the Florida Keys (Ferdie and Fourqurean 2004). Although the sediment and root tissue contained

**Table 1.** Nutrient Retention Budget for Plots with Nitrogen (N) and/or Phosphorus (P) Added at Six Sites in Florida Bay.

Treatment	Nitrogen			Phosphorus		
	C	N	NP	C	P	NP
Duck						
Sediment	61.4	140.2	145.6	2.1	56.3	50.6
<i>Thalassia</i> aboveground tissue	5.2	10.4	25.5	0.2	1.3	1.7
<i>Thalassia</i> belowground tissue	61.0	52.5	72.6	1.4	8.9	7.6
Other seagrass aboveground tissue	0.0	0.0	8.0	0.0	0.2	0.5
Other seagrass belowground tissue	0.0	0.4	19.1	0.0	0.5	2.2
Macroalgae	9.2	1.9	1.2	0.1	0.0	0.0
<i>Thalassia</i> epiphytes	0.5	0.7	1.0	0.0	0.1	0.1
Water	0.1	0.1	0.1	0.0	0.0	0.0
Total system	137.3	206.1	273.0	3.8	67.2	62.7
Net retention		68.8	135.7		63.4	58.9
Mass applied		376.3	376.3		47.3	47.3
Retention efficiency (%)		18.3	36.1		134.2	124.6
S Nest						
Sediment	43.0	49.0	127.0	1.5	38.5	49.8
<i>Thalassia</i> aboveground tissue	16.9	15.2	44.8	0.4	2.2	2.0
<i>Thalassia</i> belowground tissue	43.8	61.2	61.0	0.8	7.6	7.5
Other seagrass aboveground tissue	0.0	0.0	0.0	0.0	0.0	0.0
Other seagrass belowground tissue	0.0	0.0	0.0	0.0	0.0	0.0
Macroalgae	6.2	3.5	4.8	0.1	0.1	0.1
<i>Thalassia</i> epiphytes	1.3	1.5	1.8	0.1	0.2	0.2
Water	0.1	0.1	0.1	0.0	0.0	0.0
Total system (g)	111.4	130.4	239.5	2.9	48.6	59.5
Net retention (g)		19.1	128.2		45.8	56.6
Mass applied (g)		376.3	376.3		47.3	47.3
Retention efficiency (%)		5.1	34.1		96.8	119.8
Bob Allen						
Sediment	108.2	161.1	281.8	4.1	45.3	40.6
<i>Thalassia</i> aboveground tissue	9.9	4.4	58.0	0.2	3.1	2.5
<i>Thalassia</i> belowground tissue	45.8	21.5	124.5	1.0	17.0	11.1
Other seagrass aboveground tissue	0.0	0.0	6.2	0.0	0.3	0.2
Other seagrass belowground tissue	0.7	0.0	11.3	0.0	0.9	0.8
Macroalgae	1.6	0.0	4.8	0.0	0.1	0.2
<i>Thalassia</i> epiphytes	0.1	0.0	0.8	0.0	0.1	0.0
Water	0.1	0.1	0.1	0.0	0.0	0.0
Total system (g)	166.3	187.1	487.6	5.3	66.8	55.5
Net retention (g)		20.8	321.2		61.5	50.2
Mass applied (g)		376.3	376.3		47.3	47.3
Retention efficiency (%)		5.5	85.4		130.2	106.3
Rabbit						
Sediment	340.7	432.6	331.6	13.1	201.5	191.6
<i>Thalassia</i> aboveground tissue	57.1	69.4	69.9	2.2	3.7	4.6
<i>Thalassia</i> belowground tissue	226.8	236.9	217.3	12.0	41.2	37.1
Other seagrass aboveground tissue	0.0	0.0	3.5	0.0	0.0	0.2
Other seagrass belowground tissue	0.0	0.0	9.7	0.0	0.0	1.2
Macroalgae	0.0	0.0	0.0	0.0	0.1	0.0
<i>Thalassia</i> epiphytes	2.0	3.4	3.2	0.2	0.5	0.8
Water	0.0	0.0	0.0	0.0	0.0	0.0
Total system (g)	626.6	742.4	635.2	27.5	247.0	235.6
Net retention (g)		115.7	8.6		219.5	208.1
Mass applied (g)		376.3	376.3		47.3	47.3
Retention efficiency (%)		30.8	2.3		464.6	440.4

Table 1. continued

Treatment	Nitrogen			Phosphorus		
	C	N	NP	C	P	NP
<b>Nine Mile</b>						
Sediment	258.4	652.1	361.7	15.0	175.7	95.9
<i>Thalassia</i> aboveground tissue	43.5	84.4	53.7	1.4	3.0	3.0
<i>Thalassia</i> belowground tissue	240.1	262.4	200.7	11.6	23.8	27.4
Other seagrass aboveground tissue	0.0	0.3	5.7	0.0	0.0	0.2
Other seagrass belowground tissue	0.0	1.1	26.5	0.0	0.0	1.1
Macroalgae	0.6	0.0	0.0	0.0	0.0	0.0
<i>Thalassia</i> epiphytes	2.2	3.4	2.4	0.3	0.3	0.3
Water	0.0	0.0	0.0	0.0	0.0	0.0
Total system (g)	544.9	1003.7	650.6	28.3	202.8	127.8
Net retention (g)		458.8	105.8		174.4	99.4
Mass applied (g)		376.3	376.3		47.3	47.3
Retention efficiency (%)		121.9	28.1		369.2	210.4
<b>Sprigger</b>						
Sediment	107.8	134.3	99.2	19.0	41.1	37.5
<i>Thalassia</i> aboveground tissue	2.3	0.0	5.8	0.2	0.8	0.4
<i>Thalassia</i> belowground tissue	9.6	11.7	15.8	2.9	5.0	3.1
Other seagrass aboveground tissue	15.9	51.1	39.0	1.5	2.6	4.3
Other seagrass belowground tissue	58.1	83.9	62.5	9.3	13.3	9.2
Macroalgae	25.2	28.4	38.3	0.9	2.1	1.7
<i>Thalassia</i> epiphytes	0.7	0.2	0.2	0.1	0.1	0.0
Water	0.0	0.0	0.0	0.0	0.0	0.0
Total system (g)	219.6	309.6	260.9	33.8	65.1	56.3
Net retention (g)		89.9	41.2		31.3	22.5
Mass applied (g)		376.3	376.3		47.3	47.3
Retention efficiency (%)		23.9	11.0		66.3	47.5

Budget was calculated in April 2006 after 3.5 years of experimental fertilization. Retention efficiencies are reported in percentages; all other units are in grams within 0.25 m<sup>3</sup> compartments (50 × 50 × 100 cm) containing 200 l of water column and 50 l of sediment. Average plant and sediment values for each treatment are used in the budget.

higher absolute quantities of nitrogen than of phosphorus, percent retention of phosphorus was remarkably high, often over 100% of the P added as fertilizer over the course of our experiments. Furthermore, we likely underestimated sediment nutrient content because those measurements were collected earlier in the study. Therefore, our determinations of P content in the sediment are conservative and suggest that P retention may be even higher than we calculated. Organic acid exudates from *Thalassia* roots can extract phosphorus bound to carbonate sediments like those in Florida Bay (Long and others 2008). Therefore, the dense aggregations of *Thalassia* roots in P-enriched plots may have been able to scavenge phosphorus from pools deeper in the sediment (Holmer and others 2006). Accordingly, the highest P retention (>200%) in our study occurred in the two sites with the highest seagrass biomass (Rabbit Key and Nine Mile Bank, Figure 5).

An alternative or complimentary mechanism driving the greater than 100% phosphorus retention rates may be linked to the entrapment of

dissolved inorganic phosphorus (DIP) in the dense seagrass beds in P-enriched treatments. In carbonate environments, DIP quickly binds to particulate matter and settles to the substrate (de Kanel and Morse 1978). Further, seagrass ecosystems trap sediments and organic matter from the water column, so that on average, 50% of the organic matter deposited in seagrass beds worldwide comes from allochthonous sources (Kennedy and others 2010). These sediments contain both organic and inorganic P. Because the rate of particle trapping increases in denser seagrass beds (Eckman 1983; Boström and Bonsdorff 2000), the increased aboveground seagrass biomass in P-addition plots may have accelerated P capture from the water column.

At most sites, phosphorus retention was greater than 100%, but the westernmost site, Sprigger Bank, had relatively low (<67%) P retention. We may have underestimated sediment P content, suggesting that absolute P retention may actually be higher than 67%. Nevertheless, Sprigger Bank had markedly lower P retention relative to the

other five study sites. This is likely a consequence of the higher energy environment of Sprigger Bank, which receives higher tidal exchange and wave energy than our other sites within the protected waters of Florida Bay. This higher energy environment probably led to increased sediment resuspension and the loss of shed seagrass leaves, compared to the more quiescent inner bay sites. Similarly, Ferdie and Fourqurean (2004) found P retention was much higher in protected inshore sites compared with more exposed sites further offshore in the backreef environment of the Florida Keys. In addition, even though nutrient uptake rates tend to be higher in high velocity environments (Morris and others 2008), most of the plant biomass at the Sprigger site was macroalgae (Figure 7). Below-ground seagrass biomass, where much of the phosphorus was retained in the other sites, was quite low (Figure 5b), limiting the size of the potential P sink at that site.

In the oligotrophic Florida Bay environment, *Thalassia* is the dominant seagrass species. *Thalassia* dominance in nutrient-poor conditions is likely due to its low demand for phosphorus, relative to *Halodule* demands (Fourqurean and others 1992b). In enriched conditions, *Halodule* proliferation and subsequent decreases in *Thalassia* abundance may be attributable to light competition, where *Halodule* production of aboveground runners limited the light available to the *Thalassia* canopy underneath it (Fourqurean and others 1995).

The colonization-competition theory of community organization predicts that the initial colonizing species after a disturbance will be the inferior competitor (Tilman and Wedin 1991). Although *Halodule* is generally considered an opportunistic species that quickly colonizes following physical disturbances (Gallegos and others 1994; Robbins and Bell 2000; Sheridan 2004; Di Carlo and Kenworthy 2008), in this study, *Halodule* was slow to colonize, probably due to the small size and patchiness of a viable propagule supply. Furthermore, *Halodule* persisted in the enriched plots for an extended period, possibly because P concentrations in the sediment can remain high for decades following enrichment (Herbert and Fourqurean 2008). Changes in resource levels have been shown to facilitate species invasions in grasslands as well, where opportunistic species can capitalize on increased nutrient availability and, once established, accelerate the displacement of the species that were previously present (Siemann and Rogers 2003).

The time scale of species shifts in response to ecosystem disturbances is linked to several factors, including the supply of viable propagules (Lotze and others 2000), competitive dynamics (Fourqurean and others 1995; Rose and Dawes 1999), and resource limitation (Armitage and others 2005). In some cases, communities can recover more quickly from occasional pulsed disturbance events than from long-term, low-intensity press disturbances (Detenbeck and others 1992). In Florida Bay, however, even short-term disturbance events such as phosphorus enrichment can initiate a slow, ramped successional process that will continue over many years, possibly decades, because of the efficient retention of the limiting resource, phosphorus, in the ecosystem (Herbert and Fourqurean 2008). This extended time scale must be considered in the management and monitoring of disturbances.

## ACKNOWLEDGEMENTS

This research was funded by a grant from the Everglades National Park (ENP) under cooperative agreement 1443CA528001022 and by the Florida Coastal Everglades Long-Term Ecological Research Program funded by the US National Science Foundation (Cooperative Agreement #DEB-9910514). Water column nutrient concentrations were provided by the SERC-FIU Water Quality Monitoring Network which is supported by SFWMD/SERC Cooperative Agreement #4600000352 as well as EPA Agreement #X7-96410603-3. Doug Morrison and Bill Perry facilitated permit issuance and use of ENP facilities. We are grateful to the many people who devoted field and laboratory time to this project. Pursell Technologies Inc. and IMC Global generously donated the nitrogen and phosphorus fertilizers, respectively, for this study. This is contribution number 509 from the Southeast Environmental Research Center at FIU.

## REFERENCES

- Armitage AR, Fourqurean JW. 2009. Stable isotopes reveal complex changes in trophic relationships following nutrient addition in a coastal marine ecosystem. *Estuaries Coasts* 32:1152–64.
- Armitage AR, Frankovich TA, Heck KL Jr, Fourqurean JW. 2005. Experimental nutrient enrichment causes complex changes in seagrass, microalgae, and macroalgae community structure in Florida Bay. *Estuaries* 28:422–34.
- Bargali K. 1997. Role of light, moisture and nutrient availability in replacement of *Quercus leucotrichophora* by *Pinus roxburghii* in Central Himalaya. *J Trop For Sci* 10:262–70.
- Boström C, Bonsdorff E. 2000. Zoobenthic community establishment and habitat complexity—the importance of seagrass



- shoot-density, morphology and physical disturbance for faunal recruitment. *Mar Ecol Prog Ser* 205:123–38.
- Campbell DR, Rochefort L, Lavoie C. 2003. Determining the immigration potential of plants colonizing disturbed environments: the case of milled peatlands in Quebec. *J Appl Ecol* 40:78–91.
- de Kanel J, Morse JW. 1978. The chemistry of orthophosphate uptake from seawater on to calcite and aragonite. *Geochim Cosmochim Acta* 42:1335–40.
- Detenbeck NE, DeVore PW, Niemi GJ, Lima A. 1992. Recovery of temperate stream fish communities from disturbance: a review of case studies and synthesis of theory. *Environ Manag* 16:33–53.
- Di Carlo G, Kenworthy WJ. 2008. Evaluation of aboveground and belowground biomass recovery in physically disturbed seagrass beds. *Oecologia* 158:285–98.
- Dosch JJ, Peterson CJ, Haines BL. 2007. Seed rain during initial colonization of abandoned pastures in the premontane wet forest zone of southern Costa Rica. *J Trop Ecol* 23:151–9.
- Duarte CM. 1995. Submerged aquatic vegetation in relation to different nutrient regimes. *Ophelia* 41:87–112.
- Dunton KH. 1996. Photosynthetic production and biomass of the subtropical seagrass *Halodule wrightii* along an estuarine gradient. *Estuaries* 19:436–47.
- Eckman JE. 1983. Hydrodynamic processes affecting benthic recruitment. *Limnol Oceanogr* 28:241–57.
- Ferdie M, Fourqurean JW. 2004. Responses of seagrass communities to fertilization along a gradient of relative availability of nitrogen and phosphorus in a carbonate environment. *Limnol Oceanogr* 49:2082–94.
- Fourqurean JW, Zieman JC. 2002. Nutrient content of the seagrass *Thalassia testudinum* reveals regional patterns of relative availability of nitrogen and phosphorus in the Florida Keys USA. *Biogeochemistry* 61:229–45.
- Fourqurean JW, Zieman JC, Powell GVN. 1992a. Phosphorus limitation of primary production in Florida Bay: evidence from C:N:P ratios of the dominant seagrass *Thalassia testudinum*. *Limnol Oceanogr* 37:162–71.
- Fourqurean JW, Zieman JC, Powell GVN. 1992b. Relationships between porewater nutrients and seagrasses in a subtropical carbonate environment. *Mar Biol* 114:57–65.
- Fourqurean JW, Powell GVN, Kenworthy WJ, Zieman JC. 1995. The effects of long-term manipulation of nutrient supply on competition between the seagrasses *Thalassia testudinum* and *Halodule wrightii* in Florida Bay. *Oikos* 72:349–58.
- Fourqurean JW, Willsie A, Rose CD, Rutten LM. 2001. Spatial and temporal pattern in seagrass community composition and productivity in south Florida. *Mar Biol* 138:341–54.
- Fourqurean JW, Muth MF, Boyer JN. 2010. Epiphyte loads on seagrasses and microphytobenthos abundance are not reliable indicators of nutrient availability in oligotrophic coastal ecosystems. *Mar Poll Bull* 60:971–83.
- Frankovich TA, Fourqurean JW. 1997. Seagrass epiphyte loads along a nutrient availability gradient, Florida Bay, USA. *Mar Ecol Prog Ser* 159:37–50.
- Gallegos ME, Merino M, Rodriguez A, Marbà N, Duarte CM. 1994. Growth patterns and demography of pioneer Caribbean seagrasses *Halodule wrightii* and *Syringodium filiforme*. *Mar Ecol Prog Ser* 109:99–104.
- Grime JP. 1977. Evidence for existence of three primary strategies in plants and its relevance to ecological and evolutionary theory. *Am Nat* 111:1169–94.
- Haase J, Brandl R, Scheu S, Schädler M. 2008. Above- and belowground interactions are mediated by nutrient availability. *Ecology* 89:3072–81.
- Hauxwell J, Cebrian J, Furlong C, Valiela I. 2001. Macroalgal canopies contribute to eelgrass (*Zostera marina*) decline in temperate estuarine ecosystems. *Ecology* 82:1007–22.
- Heck KL Jr, Valentine JF, Pennock JR, Chaplin G, Spitzer PM. 2006. Effects of nutrient enrichment and grazing on shoalgrass *Halodule wrightii* and its epiphytes: results of a field experiment. *Mar Ecol Prog Ser* 326:145–56.
- Herbert DA, Fourqurean JW. 2008. Ecosystem structure and function still altered two decades after short-term fertilization of a seagrass meadow. *Ecosystems* 11:688–700.
- Holmer M, Carta C, Andersen FØ. 2006. Biogeochemical implications for phosphorus cycling in sandy and muddy rhizosphere sediments of *Zostera marina* meadows (Denmark). *Mar Ecol Prog Ser* 320:141–51.
- Hughes AR, Bando KJ, Rodriguez LF, Williams SL. 2004. Relative effects of grazers and nutrients on seagrasses: a meta-analysis approach. *Mar Ecol Prog Ser* 282:87–99.
- Jeffrey SW, Humphrey GF. 1975. New spectrophotometric equations for determining chlorophylls *a*, *b*, *c*<sub>1</sub>, and *c*<sub>2</sub> in higher plants, algae and natural phytoplankton. *Biochimie und Physiologie der Pflanzen* 167:191–4.
- Johnson NC, Rowland DL, Corkidi L, Allen EB. 2008. Plant winners and losers during grassland N-eutrophication differ in biomass allocation and mycorrhizas. *Ecology* 89:2868–78.
- Kennedy H, Beggins J, Duarte CM, Fourqurean JW, Holmer M, Marbà N, Middelburg JJ. 2010. Seagrass sediments as a global carbon sink: isotopic constraints. *Global Biogeochem Cycles* 24:GB4026. doi:10.1029/2010GB003848.
- Lake PS. 2000. Disturbance, patchiness, and diversity in streams. *J N Am Benth Soc* 19:573–92.
- Lee K-S, Dunton KH. 1999. Inorganic nitrogen acquisition in the seagrass *Thalassia testudinum*: development of a whole-plant nitrogen budget. *Limnol Oceanogr* 44:1204–15.
- Lee K-S, Dunton KH. 2000. Effects of nitrogen enrichment on biomass allocation, growth, and leaf morphology of the seagrass *Thalassia testudinum*. *Mar Ecol Prog Ser* 196:39–48.
- Long MH, McGlathery KJ, Zieman JC, Berg P. 2008. The role of organic acid exudates in liberating phosphorus from seagrass-vegetated carbonate sediments. *Limnol Oceanogr* 53:2616–26.
- Lotze HK, Worm B, Sommer U. 2000. Propagule banks, herbivory and nutrient supply control population development and dominance patterns in macroalgal blooms. *Oikos* 89:46–58.
- Martinetto P, Teichberg M, Valiela I. 2006. Coupling of estuarine benthic and pelagic food webs to land-derived nitrogen sources in Waquoit Bay, Massachusetts, USA. *Mar Ecol Prog Ser* 307:37–48.
- McGlathery KJ. 2001. Macroalgal blooms contribute to the decline of seagrass in nutrient-enriched coastal waters. *J Phycol* 37:453–6.
- MCSM. 2001. Monroe County Stormwater Management Master Plan; Vol. 1; Sect. 2.3; Pollution loads targets and analysis.
- Micheli F. 1999. Eutrophication, fisheries, and consumer-resource dynamics in marine pelagic ecosystems. *Science* 285:1396–8.
- Morris EP, Peralta G, Brun FG, van Duren L, Bouma TJ, Perez-Llorens JL. 2008. Interaction between hydrodynamics and seagrass canopy structure: spatially explicit effects on ammonium uptake rates. *Limnol Oceanogr* 53:1531–9.

- Mutchler T, Sullivan MJ, Fry B. 2004. Potential of  $^{14}\text{N}$  isotope enrichment to resolve ambiguities in coastal trophic relationships. *Mar Ecol Prog Ser* 266:27–33.
- Perez M, Duarte CM, Romero J, Sand-Jensen K, Alcoverro T. 1994. Growth plasticity in *Cymodocea nodosa* stands: the importance of nutrient supply. *Aquat Bot* 47:249–64.
- Pickett STA, Kolasa J, Armesto JJ, Collins SL. 1989. The ecological concept of disturbance and its expression at various hierarchical levels. *Oikos* 54:129–36.
- Powell GVN, Kenworthy WJ, Fourqurean JW. 1989. Experimental evidence for nutrient limitation of seagrass growth in a tropical estuary with restricted circulation. *Bull Mar Sci* 44:324–40.
- Reed DC, Raimondi PT, Carr MH, Goldwasser L. 2000. The role of dispersal and disturbance in determining spatial heterogeneity in sedentary organisms. *Ecology* 81:2011–26.
- Robbins BD, Bell SS. 2000. Dynamics of a subtidal seagrass landscape: seasonal and annual change in relation to water depth. *Ecology* 81:1193–205.
- Rose CD, Dawes CJ. 1999. Effects of community structure on the seagrass *Thalassia testudinum*. *Mar Ecol Prog Ser* 184:83–95.
- Sheridan P. 2004. Recovery of floral and faunal communities after placement of dredged material on seagrasses in Laguna Madre, Texas. *Estuar Coast Shelf Sci* 59:441–58.
- Short FT. 1983. The seagrass, *Zostera marina* L.: plant morphology and bed structure in relation to sediment ammonium in Izembek Lagoon, Alaska. *Aquat Bot* 16:149–61.
- Siemann E, Rogers WE. 2003. Changes in light and nitrogen availability under pioneer trees may indirectly facilitate tree invasions of grasslands. *J Ecol* 91:923–31.
- Tilman D. 1987. Secondary succession and the pattern of plant dominance along experimental nitrogen gradients. *Ecol Monogr* 57:189–214.
- Tilman D, Wedin D. 1991. Dynamics of nitrogen competition between successional grasses. *Ecology* 72:1038–49.
- Valiela I, McClelland J, Hauxwell J, Behr PJ, Hersh D, Foreman K. 1997. Macroalgal blooms in shallow estuaries: controls and ecophysiological and ecosystem consequences. *Limnol Oceanogr* 42:1105–18.
- Vitousek PM, Aber JD, Howarth RW, Likens GE, Matson PA, Schindler DW, Schlesinger WH, Tilman DG. 1997. Human alteration of the global nitrogen cycle: sources and consequences. *Ecol Appl* 7:737–50.
- Welschmeyer NA. 1994. Fluorometric analysis of chlorophyll *a* in the presence of chlorophyll *b* and pheopigments. *Limnol Oceanogr* 39:1985–92.
- Zieman JC, Fourqurean JW, Iverson RL. 1989. Distribution, abundance and productivity of seagrasses and macroalgae in Florida Bay. *Bull Mar Sci* 44:292–311.